







DINOSAURIA – distinguishing characters


- advanced mesotarsal ankle
- limbs beneath the body, upright posture
- (a sacrum composed of three or more fused vertebrae)
- (open hip socket, the acetabulum)
- asymmetric hand with two smaller outer digits
- elongate deltopectoral crest on the upper arm bone




according to Brusatte (2011)

- temporal musculature
(mandibular adductors enlarged)
- epipophyses on cervical vertebrae
- elongate deltopectoral crest on humerus
- open acetabulum
- asymmetrical fourth trochanter
- bifurcating posterior jugal process










Marginocephalia	Ornithopoda	Stegosauria	Arkylosauria	Prosauropoda	Sauropoda	Carnosauria etc.	Birds
Cerapoda		Thyreophora		Sauropodomorpha		Theropoda	
Ornithischia 				Saurischia 			

DINOSAURIA




Coelophysis

Ornithischia




Ornithischian characters

- "bird hip" (pubis points backward)
- predentary bone; premaxillae expanded and toothless
- small antorbital fenestrae
- blunt or leaf shaped teeth
- ossified tendons along parts of spinal column
- five or more sacral vertebrae
- herbivorous (omnivorous)
- since the late Triassic




Ornithischia – group 1



ORNITHISCHIA


Thyreophora (“shield-bearers - *sköldbärare*”)



- Rows of dermal ossification (armored dinosaurs)



Thyreophora – stem forms – Stegosauria / Ankylosauria




Thyreophora – stem forms – Stegosauria / Ankylosauria


Scutellosaurus

- Relatively small (1 m long), biped
- armored with several hundreds scutes (or osteoderms)
- only two partial skeletons known
- Early Jurassic, North America





Thyreophora – **stem forms** – Stegosauria / Ankylosauria






Thyreophora – stem forms – Stegosauria / Ankylosauria

Scelidosaurus

- quadruped (ca. 4 m long)
- heavily armored
- Early Jurassic, Europe and North America






Thyreophora – **stem forms** – Stegosauria / Ankylosauria



Scelidosaurus

Thyreophora – stem forms – Stegosauria / Ankylosauria





Thyreophora – stem forms – **Stegosauria** / Ankylosauria

Stegosauria (Jurassic to Cretaceous)

("roof-lizards, plated lizards - taködlor")

- quadrupedal; legs with hoof-like toes
- long, narrow heads and a horn-covered toothless beak
- head relatively small to body size
- weak teeth
- row(s) of osteoderms, which develop into plates and spines along the back and tail
- long neck and tail



small antorbital fenestrae, toothless predentary bone




Thyreophora – stem forms – **Stegosauria** / Ankylosauria

Stegosauria

- Appear during the Middle Jurassic
- Diversify during the Middle and Late Jurassic
- Decline during the Cretaceous and go extinct at the end of the Cretaceous (not because of their small brain)


THE FAR SIDE

By GARY LARSON




Stegosaurus Marsh, 1879

the best known, and also most famous stegosaur
... and the trouble with the plates (osteoderms)



Stegosaurus and the second brain



large canal in the spinal cord (20 times larger than the brain)
→ glycogen body ??

Thyreophora – stem forms – **Stegosauria** / Ankylosauria

Ode to *Stegosaurus* by B. L. Taylor (1912)

“[...]

If one brain found the pressure strong

It passed a few ideas along.


If something slipped his forward mind

'Twas rescued by the one behind.

And if in error he was caught

He had a saving afterthought.

[...]”





Thyreophora – stem forms – Stegosauria / **Ankylosauria**

Ankylosauria

(“fused lizards” – “*pansardinosaurier*”)

- herbivorous
- quadruped – up to 9 m
- long body with relatively short legs
- heavily armored with osteoderms
- cranial ornamentation, reduction of cranial fenestrae
- wide body
- weak jaws with small teeth
- Often with club at the end of tail
- typical for the Cretaceous (since Late Jurassic)





Thyreophora — stem forms — Stegosauria / **Ankylosauria**


Ankylosauria and their armor

JURASSIC ANKYLOSAURS



Scelidosaurus

CRETACEOUS ANKYLOSAURS



Nodosaurus

Polacanthus

Minmi

Ankylosaurus


Euoplocephalus

Pinacosaurus

- armor is rarely preserved *in situ*
- dermal plates most likely arranged in rows (longitudinally and/or transversely)

Thyreophora — stem forms — Stegosauria / **Ankylosauria**

Gastonia - Early Cretaceous, Moab, Utah




Thyreophora — stem forms — Stegosauria / **Ankylosauria**

Reduction of cranial fenestrae




Thyreophora — stem forms — Stegosauria / **Ankylosauria**

Ankylosaur synsacrum




Thyreophora — stem forms — Stegosauria / Ankylosauria



Thyreophora – stem forms – Stegosauria / **Ankylosauria**

Nodosauridae


- narrow skull without excessive armor or horns
- presence of spines as armor
- lack of tail club
- open lateral temporal fenestrae



Polacanthus

Ankylosauridae

- wide armored head with horns
- few or no spines as armor
- presence of tail club
- closed fenestrae (i.e., covered with bony armor)



Euoplocephalus


Euoplocephalus



Thyreophora — stem forms — Stegosauria / **Ankylosauria**



Ornithischia – group 2




ORNITHISCHIA

Cerapoda ("horned feet")

shared character:


→ thicker layer of enamel on the inside of their lower teeth



Cerapoda – Marginocephalia + Ornithopoda


Ornithopoda - (“bird feet”)

- bipedal or quadrupedal
- no body armor
- since early Jurassic
- diverse group
 - heterodontosaurids
 - iguanodontids
 - duck-billed dinosaurs
- ventrally offset tooth rows
- low located lower jaw joint



Cerapoda – Ornithopoda – Heterodontosauridae – Iguanodontidae – Hadrosauridae



ORNITHOPODA



Cerapoda – Ornithopoda – Heterodontosauridae – Iguanodontidae – Hadrosauridae

Hypsilophodonts




- Mostly small (1-3 m long), obligate bipeds
- Probably fast runners
- Earliest forms in Middle Jurassic, persist until end of Cretaceous
- Most famous representative is the hypsilophodontid *Hypsilophodon*



special way of grinding food → probably a key to their success


Iguanodontidae

- Almost all are **larger** than "hypsilophodonts"
- strong forelimbs, and were probably **facultative bipeds**
 - bipedal when running and feeding on trees
 - quadrupedal while walking and browsing on low vegetation
- Characterized by **loss of premaxillary teeth**
- **spike-like thumb** present in many iguanodontids
- Since Middle Jurassic, but become more common in Late Jurassic and Early Cretaceous
- *Camptosaurus* (Late Jurassic, North America)
Tenontosaurus (Early Cretaceous, North America)
and *Iguanodon* (Early Cretaceous, Europe)



IGUANODON

1 meter
3 feet



Iguanodon (ig-WHA-noh-don) "iguana tooth"
Period: Late Jurassic and Early Cretaceous
Length: 30 feet (9 meters)
Location: Africa, Asia, Australia, Europe, and North America

A massive herbivore with a horny beak and cheek teeth for grinding vegetation. Its hands had distinctive hoofed fingers and spiked thumbs.

© 2005 Encyclopædia Britannica, Inc.

Hadrosauridae - "duck-billed dinosaurs"

- Extremely common in Late Cretaceous of North America, also known from Asia, Europe, South America
- one of the best known dinosaur group (from eggs through adults)
- lived in herds and had nesting colonies
- **End of snout flares outward** to form "duck bill"
 - increase in number of teeth
 - Teeth organized into **dental battery**
a continuous grinding surface
 - most complex and sophisticated chewing of any reptile
 - **Manual digit I lost**
 - **Metacarpals II-IV very long and slender:**
 - probably quadruped most of the time





Cerapoda — Ornithopoda — Hypsilophodonts — Iguanodontidae — **Hadrosauridae**



Hadrosaur Teeth
Some hadrosaurs had more than 1200 teeth packed together along the sides of their mouths. Banks of closely-packed teeth enabled them to chop tough plant material into fine, easily-digested pieces.

Hadrosaur Dentary
Dinosaur Provincial Park, Alberta

Hadrosaur Tooth
Dinosaur Provincial Park, Alberta

Hadrosaur Jaw
Hadrosaurs had broad duck-like snouts. They were plant eaters, and their teeth were worn down

Cerapoda — Ornithopoda — Hypsilophodonts — Iguanodontidae — **Hadrosauridae**

Hadrosauridae - "duck-billed dinosaurs"




Two major clades:

Hadrosaurinae

→ enlarged external nares
and broader snouts




Lambeosaurinae

→ hollow crests on top of head



Possible uses of lambeosaurine crests?

- Increase surface area for smelling sensors
- Sound generation 
- Visual display
- Trapping moisture in exhaling breath to keep lungs from drying out
- Or a combination
- are different in every species
- probably display also sexual dimorphism



Cerapoda — Ornithopoda — Hypsilophodonts — Iguanodontidae — **Hadrosauridae**




How to speak dinosaur

- [\(http://www.courtney-brown.net/\)](http://www.courtney-brown.net/)
- <http://prix2015.aec.at/prixwinner/15549/>

ORNITHISCHIA

Marginocephalia (“fringed heads” - “*kragförsedda*”)


- all herbivores
- bi- or quadrupedal
- characterized by a bony ridge or frill on the back of the skull
- evolved during the Jurassic
- especially Upper Cretaceous
- Ceratopsia + Pachycephalosauria



The five dinosaur faunas

	Fauna	Theropoda	Sauropodomorpha	Ornithopoda	Thyreophora	Marginocephalia
Cretaceous	Late	5 Carnosauria: (Tyrannosauridae) Coelurosauria: (Ornithomimosauridae) (Dromaeosauridae) (Oviraptorosauridae)		Hadrosauridae	Ankylosauria: Ankylosauridae	Neoceratopsians: Protoceratopsidae Ceratopsidae Pachycephalosauria
	Early	4	Sauropoda: Small Sauropods (Titanosaurids)	Large Ornithopods: Iguanodontidae	Ankylosauria: Nodosauridae Stegosauria: A few Stegosaurids	Psittacosaur
Jurassic	Late	3 Carnosauria: (Allosauridae)	Sauropoda: Large Sauropods	Medium Ornithopods: Hypsilophodontidae Iguanodontidae	Stegosauria: Large Stegosaurids	
	Early	2	Sauropoda: Small Sauropods (Cetiosaurids)	Small Ornithopods: Hypsilophodontidae	Stegosauria: Small Stegosaurids Primitive Thyreophorans	
Triassic	Late	1 Ceratosauria Also Primitive Saurischians	Prosauropods	Primitive Ornithischians		

Marginocephalia: evolve during the Middle Jurassic,
but become common during the Cretaceous




Cerapoda – Marginocephalia – **Pachycephalosauria** – Ceratopsia


Pachycephalosauria

("thick headed lizards"- "*benskallar*")


- Since Cretaceous (esp. North America & Asia)
 - massive, thick skull roof (fusing and thickening of frontal & parietal bones)
- typically no supratemporal openings



Drakorex



Stygimoloch



Pachycephalosaurus


Pachycephalosauria

- long hindlimbs, short forelimbs
- long tail with vertebrae tightly joined by ossified tendons
 - much like ornithomimids
- long ilium bone
- six to eight sacral vertebrae
- head placed at an angle to the vertebral column
- bipedal
- Sexual dimorphism
- similar to the ornithomimids




Head butting?

- thick skull to protect brain
- shortened neck
- strengthened vertebral column
- inclination of the skull



Cerapoda — Marginocephalia — **Pachycephalosauria** — Ceratopsia



but ...



Microstructure showed:

- internal structure of the pachycephalosaur dome is a dynamic tissue, fast growing bone (ontogenetic stage of dome growth)
- dense layer of Sharpey's fibers indicates external covering



Photos courtesy John Horner / Museum of the Rockies (above); M. Goodwin (left)

Goodwin & Horner, 2004




Cerapoda — Marginocephalia — **Pachycephalosauria** — Ceratopsia



Cerapoda – Marginocephalia – **Pachycephalosauria** – Ceratopsia




Cerapoda – Marginocephalia – **Pachycephalosauria** – Ceratopsia






Ceratopsia ("horned faces" - "*hornansikte*")

- generally quadrupedal
- triangular head
- **rostral bone**
- skull with narrow beak
- advanced forms with shelf of bone (and horn)
- Cretaceous of Asia and North America




Psittacosaurus "the parrot dinosaur"


- ancestral to the Ceratopsia
 - similar to ornithomimids
 - most rudimentary frill
 - rostrum, beak, short snout, high position of the nostrils
- primitive ceratopsian



Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia**





- *Psittacosaurus* (early Cretaceous of China)
- bristle-like structures on the proximal part of tail



Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **psittacosaurids**

Protoceratops and the protoceratopsids

- still relatively small (1-2.5 m)
- intermediate between psittacosaurus and true ceratopsids
- forelimbs become equal in length with the hind limbs
 - quadrupedal
- Late Cretaceous of North America and Asia (esp. Mongolia and China)
- no horns, rather small frill
- small nostrils
 - more primitive



Cerapoda — Marginocephalia — Pachycephalosauria — **Ceratopsia** — **protoceratopsids**

The Griffin Myth



ca. 1260, France



ca. 375-350 BC, Eritrea

Griffin: a lion with an eagle head that lies eggs

first described by Greek writers 675 BC

→ finds of protoceratopsids in Mongolia by
Scythian nomads



Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **protoceratopsids**



Yinlong downsi
Oxfordian, China



Chaoyangsaurus youngi
Tithonian, China



Auroraceratops rugosus
Barremian, China



Liaoceratops yanzigouensis
Barremian, China



Psittacosaurus mongoliensis
Aptian, Mongolia & China



Hongshanosaurus houi
Barremian, China



Archaeoceratops oshimai
Aptian, China



Yamaceratops dongobiensis
Albian, Mongolia



Psittacosaurus sibiricus
Albian, Russia



Zuniceratops christopheri
Turonian, New Mexico



Graciliceratops mongoliensis
Santonian-Campanian,
Mongolia



Protoceratops andrewsi
Santonian-Campanian,
Mongolia & China



Cerasinops hodgskissi
Campanian, Montana



Udanoceratops tschizhovi
Campanian, Mongolia



Bagaceratops rozhdestvenskyi
Campanian, Mongolia



Leptoceratops gracilis
Maastrichtian, Wyoming &
Alberta



Montanoceratops cerorhynchus
Maastrichtian, Montana






Prenoceratops pieganensis
Campanian, Montana

1 m



Ceratopsidae


- all quadrupedal
- enlarged frill
 - skull up to 2.4 m long (parietal and squamosal bones)
- variety of horns
- **enlarged nares**
- 4 to 8 m long
- only Late Cretaceous of North America




Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**

Two groups: Chasmosaurinae and Centrosaurinae


long-frilled, long-faced vs. short-frilled, short faced forms




Triceratops



Pentaceratops



Centrosaurus




Styracosaurus

- large brow horns
- enlarged and long frill
- enlarged rostral bone and thickening of the snout bones

- large nasal horn
- short brow horns
- hooks and spines on frill margin


Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**






Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**



Cerapoda — Marginocephalia — Pachycephalosauria — **Ceratopsia** — **ceratopsids**




Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia – ceratopsids**




Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**





Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**




Function of the frill ??



- muscle attachment
- reflects gender
- ritualized combat and display
- establishment of hierarchies
- protection of the neck
- thermoregulation

Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**



Function of the horns ??



- predator defense
- intimidate predator
- mate recognition
- intraspecific competition

Cerapoda — Marginocephalia — Pachycephalosauria — **Ceratopsia** — **ceratopsids**

Sexual dimorphism



Chasmosaurus

Cerapoda – Marginocephalia – Pachycephalosauria – **Ceratopsia** – **ceratopsids**